

FUMAGRI EFFISAFE



Pig &
Poultry
Farming

A broad spectrum
of activity disinfectant
for use at livestock
premises, equipment,
foot bath, animal
transport trucks and
wheel disinfection.




FUMAGRI EFFISAFE



ADVANTAGES

- Disinfectant based on two active ingredients
- Neutral solution at 1% dilution
- Kind to many surfaces
- Bactericidal, yeasticidal, fungicidal, mycobactericidal and virucidal disinfectant for livestock premises, animal breeding equipment and materials in farms.
- Bactericidal, yeasticidal and virucidal disinfectant for animal transport vehicles, including wheels, boots in bootbath



Surface / Equipment Disinfection

Pre-clean surfaces and equipment to be disinfected.

Prepare a dilution of FUMAGRI EFFISAFE and apply the solution with 300ml per m² (depending on the surface) by low pressure spraying, foaming* or by soaking at a minimum of 10°C:

- 1% bactericidal, yeasticidal, virucidal / ≥ 30 min.
- 3% fungicidal / ≥ 30 min.
- 1% mycobactericidal and 2% fungicidal / ≥ 60 min.

Use the disinfecting solution within 24 hours after preparation.

Foot, wheel or animal transport truck disinfection:

Prepare a dilution and apply by low pressure spraying or foaming* on surfaces for animal transport trucks and wheels or apply as a boot or wheel bath by soaking.

- 2% Bactericidal, yeasticidal, virucidal / >= 5 min.

Renew the disinfection bath on a daily base.

* For foaming gun, use a 25% dilution mother solution in the tank.

BACTERICIDAL EFFICACY

- EN14349 / 10°C / low soiling:
 - 2%, 30 min.: *Campylobacter jejuni*
 - 2%, 5 min.: *Enterococcus hirae*, *Proteus hauseri*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*
- EN1656 / 10°C / low soiling
 - 0.5%, 1 min.: *Proteus vulgaris*
 - 1%, 1 min.: *Staphylococcus aureus*
 - 2%, 1 min.: *Enterococcus hirae*, *Pseudomonas aeruginosa*
 - 1%, 5 min.: *Enterococcus hirae*, *Proteus hauseri*, *Pseudomonas aeruginosa*, *Staphylococcus Aureus*
 - 0.2%, 30 min.: *Proteus hauseri*
 - 0.5%, 30 min.: *Enterococcus hirae*,
 - 1%, 30 min.: *Pseudomonas aeruginosa*, *Staphylococcus aureus*

FUNGICIDAL EFFICACY

- EN16438 / 10°C / low soiling
 - 1%, 60 min.: *Aspergillus fumigatus*
 - 2%, 60 min.: *Aspergillus brasiliensis*
 - 3%, 30 min.: *Aspergillus brasiliensis*
- EN1657 / 10°C / low soiling
 - 2%, 30 min. *Aspergillus brasiliensis*

VIRUCIDAL EFFICACY

- EN14675 / 10 °C / low soiling
 - 1.5%, 1 min.: *Bovine Enterovirus type 1 (ECBO)*
 - 0.5%, 30 min.: *Bovine Enterovirus type 1 (ECBO)*
 - 0.75%, 30 min.: *Classical Swine Fever's disease virus*
 - 1%, 1 min.: *African swine fever virus (ASF)*
- EN14675 / 10 °C / high soiling
 - 1%, 5min.: *Bovine Enterovirus type 1 (ECBO)*
 - 2%, 1 min.: *African swine fever virus (ASF)*

YEASTICIDAL EFFICACY

- EN16438 / 10°C / low soiling
 - 2%, 1 min.: *Candida albicans*
 - 1%, 5 min. & 30 min.: *Candida albicans*
- EN 1657 / 10°C / Low soiling / 2% / 1 min
- EN 1657 / 10°C / Low soiling / 1% / 30 min

MYCOBACTERICIDAL EFFICACY

- EN14204 / 10°C / low soiling / 1% / 60 min.: *Mycobacterium avium*

Active substances:

Chlorocresol & OPP (Orthophenylphenol)

pH at 10 g/l:

7.39±0.5

Storage:

Keep in a clean, cool and well-ventilated place away from sources of heat and intense light.

Shelf Life:

2 years from date of manufacture

For complete information on products and availability according to local regulation please contact: Poultryandpigs@kersia-group.com
Dangerous. Respect precautions of use. Disinfectants are regulatory biocides. They have guarantees of efficiency and protection of human, animals and the environment. Use biocides safely. Before use read the label and product information.